

Atomic data from the Iron Project

LIX. New radiative transition probabilities for Fe IV including fine structure

S. N. Nahar and A. K. Pradhan

Department of Astronomy, The Ohio State University, Columbus, OH 43210, USA
e-mail: nahar@astronomy.ohio-state.edu *

the date of receipt and acceptance should be inserted later

Abstract. We present new calculations for transition probabilities of Fe IV, with much more extensive datasets than heretofore available. The large-scale close coupling R-matrix calculations yield 1,798 LS bound states with $n \leq 11$ and $l \leq 9$, and corresponding transition probabilities for 138,121 dipole allowed transitions in the form of oscillator strengths f , line strengths S , and A-coefficients for a variety of applications. This represents the largest R-matrix dataset in LS coupling for any ion under either the Opacity Project or the Iron Project. Through algebraic transformation of the LS multiplets, a total of 712,120 dipole allowed fine structure transitions for Fe IV are obtained. Observed transition energies are used together with the energy independent line strengths to derive the f - and the A-values; the adopted algorithm used calculated energies for the remainder. Present results show significantly better accuracy for the important low-lying states than previous calculations. Monochromatic and mean opacities for Fe IV are computed and compared with those obtained using the Opacity Project data. We find differences which could have important consequences for several astrophysical applications involving low ionization stages of iron.

Key words. atomic data – radiative transition probabilities – fine structure transitions – opacities – Ti-like ions – ultraviolet radiation sources

1. Introduction

It is well known that low ionization stage of iron are of great interest for astrophysical spectroscopy and models of stellar atmospheres (Hubeny *et al.* 2003), novae (McKenna *et al.* 1997), photoionized H II regions (Dopita and Sutherland 2003, Rubin *et al.* 1997), and active galactic nuclei (Sigut and Pradhan 2003, Sigut *et al.* 2004). The ion Fe IV is of particular interest since its ionization potential is ~ 54.8 eV, nearly equal that of He II at 54.4 eV. Thus, for example, in a photoionized H II region such as the great diffuse nebula in Orion, stellar ionizing flux is effectively shielded by Helium to prevent extensive ionization of Fe ions beyond Fe IV. Whereas hot central stars of high-excitation planetary nebulae may well ionize beyond Fe IV (e.g. Hyung and Aller 1998), ordinary main sequence O, B stars have a 54 eV 'cut-off' that attenuates the ionizing flux. Another example is the anomalous Fe II emission from active galactic nuclei. Non-local-Thermodynamic-Equilibrium (NLTE) spectral models need to consider the first four ionization stages, Fe I - IV, with Fe IV as the dominate stage in the fully ionized zone (e.g. Sigut *et al.* 2004 and references therein). In previous

studies at Ohio State under the Iron Project, we have reported a large amount of radiative and collisional data for the ionization stages Fe I-VI, including fine structure transition probabilities for Fe II and Fe III (references at: www.astronomy.ohio-state.edu/~pradhan). The aim of this paper is to report similarly new calculations with improved accuracy for Fe IV including fine structure that is often required for spectral diagnostics and modeling.

However, study of Fe IV is difficult and complexities arise due to its electronic structure with 5 electrons in the half-filled open 3d shell (ground configuration $1s^2 2s^2 2p^6 3s^2 3p^6 3d^5$); electron correlation effects for the many excited states are difficult to represent. Prior to the Opacity Project (OP, 1995, 1996), investigations on radiative processes for Fe IV were carried out through atomic structure calculations. These include radiative decay rates for forbidden transitions by Garstang (1958), which are available from the evaluated compilation by the National Institute for Standards and Technology (NIST: www.nist.gov) and Rubin and Froese-Fisher (1998). Fawcett (1989) has computed the data for many allowed transitions using a semi-empirical method due to Cowan.

Ab initio calculations in the close coupling approximation using R-matrix method for the radiative processes were carried out under the Opacity Project (OP, 1995, 1996) by Sawey and Berrington (1992), and later under the Iron Project (IP, Hummer *et al.* 1993) by Bautista and Pradhan (1997).

Send offprint requests to: S. N. Nahar

* Complete electronic data tables of energies and transition probabilities are available from the CDS via anonymous ftp to cdsarc.u-strasbg.fr (130.79.128.5) or via <http://cdsweb.u-strasbg.fr/Abstract.html>

Further R-matrix calculations were reported for the total and state-specific electron-ion recombination rate coefficients using the unified treatment by Nahar, Bautista, and Pradhan (1998). Sawey and Berrington (1992) produced radiative data for photoionization and oscillator strengths for three symmetries: sextets, quartets and doublets; these data are accessible through the OP database, TOPbase (Cunto *et al.* 1993). Later work of Bautista and Pradhan (1997) took more accurate account of electron-electron correlation effects, showing important resonant features in cross sections missing in the former investigation by Sawey and Berrington (1992). However, owing to computational constraints, Bautista and Pradhan (1997) entirely omitted the doublet symmetry which has the maximum number of bound states. All previous R-matrix results are in LS coupling and consider only the dipole allowed LS multiplets; no fine structure splitting has been taken into account.

Present work reports a more complete set of oscillator strengths and radiative decay rates of Fe IV for both the LS multiplets and allowed fine structure transition. Similar results have been presented earlier for Fe II (Nahar 1994), Fe III (Nahar and Pradhan 1996), and for some other ions. Present calculations consider correlation effects accurately, as in Bautista and Pradhan (1997), and bound states of all symmetries, sextets, quartets, and doublets. At the present time it is computationally prohibitive to carry out full Breit-Pauli R-matrix calculations for fine structure transition probabilities including relativistic effects in an *ab initio* manner. Among the low ionization stages of iron, Fe I - V, there has been only one such previous effort, for Fe V (Nahar and Pradhan 2000, Nahar *et al.* 2000). challenging computationally. While efforts are under way to resolve the computationally challenging problems in relativistic calculations, which are about an order of magnitude more intensive, there is considerable need for fine structure data. We employ an algebraic recoupling algorithm to obtain the fine structure transition probabilities for Fe IV, using observed energies where available, as in previous IP works on Fe II and Fe III.

Finally, we employ the new radiative dataset to compute monochromatic and mean opacities for Fe IV, both as a test and comparison with the earlier OP work, as well as to investigate possible effect on the astrophysically important problem of iron opacities.

2. Theory

The R-matrix method as applied to the OP and IP work enables the consideration of many excited bound states and transitions, typically for all states with $n \leq 10$. General details of the procedures employed have been described in earlier papers, such as for Fe II (Nahar 1993), for obtaining oscillator strengths (f) and radiative decay rates (A) for dipole allowed fine structure transitions ($\Delta J = 0, \pm 1$ with $\Delta S = 0$, even parity $\leftrightarrow$ odd parity) by algebraic transformation of LS multiplets. The methodology of the close coupling (CC) approximation employing the R-matrix method for radiative processes is discussed in the OP papers by Seaton (1987) and Berrington *et al.* (1987). The wavefunction $\Psi(E)$ for the $(N+1)$ electron system with total spin and orbital angular momenta symmetry $SL\pi$ or total angu-

lar momentum symmetry $J\pi$ is expanded in terms of ‘frozen’ N -electron target ion functions χ_i and vector coupled collision electrons θ_i ,

$$\Psi_E(e + ion) = \mathcal{A} \sum_i \chi_i(ion) \theta_i + \sum_j c_j \Phi_j(e + ion), \quad (1)$$

in some specific state $S_i L_i \pi_i$ or level $J_i \pi_i$, index i marking channels $S_i L_i (J_i) \pi_i$ $k_i^2 \ell_i (SL\pi$ or $J\pi)$ with energy k_i^2 of the colliding electron described by θ . $\mathcal{A}$ is the antisymmetrization operator. The second sum expands correlation functions Φ_j as products with $N+1$ bound orbital functions that (a) compensate for the orthogonality conditions between the continuum and the bound orbitals, and (b) represent additional short-range correlation that is often of crucial importance in scattering and radiative CC calculations for each $SL\pi$. c_j 's are variational coefficients.

R-matrix solutions of coupled equations to total symmetries LS states for $(N+1)$ -electron system,

$$H_{N+1} \Psi = E \Psi, \quad (2)$$

are discrete bound states Ψ_B for $E < 0$. Seaton (1985) describes the matching procedure for inner region and outer region wavefunctions at the R-matrix boundary which enables calculations of all required energy levels of the Hamiltonian matrix.

For a bound-bound transition from an initial state i to a final state j , the line strength, S , is defined as

$$S = |\langle \Psi_j | \mathbf{D} | \Psi_i \rangle|^2, \quad (3)$$

in atomic units (a.u.), where the dipole operator $\mathbf{D}$ is

$$\mathbf{D} = \sum_n \mathbf{r}_n, \quad (4)$$

in the length form, summed over the total number of electrons in the ion, and Ψ_i and Ψ_j are the initial and the final wavefunctions respectively. The oscillator strength f_{ij} (f -value) can be obtained from line strength S in atomic units ss

$$S = (3g_i/E_{ij}) f_{ij}. \quad (5)$$

where E_{ij} is the transition energy in rydbergs and g_i is the statistical weight factor of the initial state, with $g_i = (2S_i+1)(2L_i+1)$ for a LS multiplet, and $(2J_i+1)$ for a fine-structure transition. The radiative transition probability, A_{ji} (Einstein's A-coefficient or the radiative decay rate), can be obtained from the oscillator strength as

$$A_{ji}(a.u.) = \frac{1}{2} \alpha^3 \frac{g_i}{g_j} E_{ij}^2 f_{ij}, \quad A_{ji}(s^{-1}) = \frac{A_{ji}(a.u.)}{\tau_0}, \quad (6)$$

where α is the fine-structure constant and $\tau_0 = 2.4191 \times 10^{-17} s$ is the atomic unit of time.

The fine-structure components of an LS multiplets can be obtained through algebraic transformations of either line strength or directly from the oscillator strength (e.g. Nahar 1995). The line strength, which is independent of the transition energy E_{ij} , is a better choice. As the observed transition energies are determined more accurately than the calculated ones, use of the former with the line strengths can provide more accurate f_{ij} and A_{ji} . Hence, present fine-structure components are

obtained from S values whenever observed energies are available, and from f -values when calculated E_{ij} are used.

The fine-structure line strengths, S_{JJ} , are obtained as

$$S_{JJ} = C(J_i, J_j) S_{LS} / [(2S_i + 1)(2L_i + 1)(2L_j + 1)], \quad (7)$$

for the allowed transitions ($\Delta J = 0, \pm 1$). S_i is the spin which is the same as S_j . The values of the coefficients $C(J_i, J_j)$ can be found in Allen (1976). The S_{JJ} values satisfy the condition

$$S_{LS} = \sum_J S_{JJ}. \quad (8)$$

The fine-structure f -values, f_{JJ} , can be obtained directly from f_{LS} as (Seaton et al 1994):

$$f_{JJ}(n_j S_i L_j J_j, n_i S_i L_i J_i) = f_{LS}(n_j S_i L_j, n_i S_i L_i) (2J_j + 1)(2L_i + 1) \times W^2(L_j L_i J_j J_i; 1S_i), \quad (9)$$

where $W(L_j L_i J_j J_i; 1S_i)$ is a Racah coefficient. The above components also satisfy the sum rule

$$\sum_{J_i J_j} (2J_i + 1) f_{JJ}(n_j S_i L_j J_j, n_i S_i L_i J_i) = f_{LS}(n_j S_i L_j, n_i S_i L_i) \times (2S_i + 1)(2L_i + 1). \quad (10)$$

The above form for fine structure components is used when one or both LS terms of the transition are unobserved or when not all the fine structure levels of an LS term are observed, and for transitions between high angular momentum states, such as higher than $H \leftrightarrow I$ for which Allen's coefficients are not available.

The lifetime of a level can be computed as

$$\tau = \frac{1}{\sum_i A_{ki}} \quad (11)$$

where the sum is the total radiative transition probability for level k , i. e.

$$g_i A_{ki}^{E1} = 2.6774 \times 10^9 \text{ s}^{-1} (E_i - E_k)^3 S^{E1}(i, k) \quad (12)$$

(the observed rate) in the electric dipole case.

The opacity is a measure of radiation transport through matter. The monochromatic opacity is a function of the photon frequency and depends mainly on the detailed atomic data: bound-bound oscillator strengths and bound-free photoionization cross sections. The quantity of practical interest in stellar models is the Rosseland mean opacity, $\kappa_R(T, \rho)$, defined as (Seaton et al. 1994)

$$\frac{1}{\kappa_R} = \int_0^\infty \frac{1}{\kappa_\nu} g(u) du, \quad g(u) = \frac{15}{4\pi^4} u^4 e^{-u} (1 - e^{-u})^{-2}, \quad (13)$$

where $g(u)$ is the Planck weighting function and $u = h\nu/kT$. Monochromatic opacities κ_ν depends primarily on oscillator strengths as

$$\kappa_\nu(i \rightarrow j) = \frac{\pi e^2}{mc} N_i f(i, j) \phi_\nu \quad (14)$$

where N_i = ion density in state i , ϕ_ν is a profile factor, and on photoionization cross sections, σ_{PI} , as

$$\kappa_\nu = N_i \sigma_{PI}(\nu). \quad (15)$$

We compute the monochromatic and the mean opacities as a measure of the differences between the present dataset for the bound-bound transitions and the earlier OP data for Fe IV.

Table 1. Target terms and relative energies (in Ry) of Fe V in the wavefunction expansion of Fe IV.

	Term	E_o/Ry
1	$3d^4 \ ^5D$	0.000000
2	$3d^4 \ ^3H$	0.230232
3	$3d^4 \ ^3P$	0.234211
4	$3d^4 \ ^3F$	0.244942
5	$3d^4 \ ^3G$	0.274991
6	$3d^4 \ ^1G$	0.333399
7	$3d^4 \ ^3D$	0.334726
8	$3d^4 \ ^1I$	0.341832
9	$3d^4 \ ^1S$	0.361166
10	$3d^4 \ ^1D$	0.421837
11	$3d^4 \ ^1F$	0.480536
12	$3d^4 \ ^3F$	0.567719
13	$3d^4 \ ^3P$	0.568463
14	$3d^4 \ ^1G$	0.649554
15	$3d^4 \ ^1D$	0.855063
16	$3d^4 \ ^1S$	1.103820

3. Computations

The R-matrix computations for Fe IV are carried out using a wavefunction expansion of 16 terms of target Fe V belonging to the ground configuration, $3d^4$, as given in Table 1; the energies used are the measured values available from the NIST compilation at www.nist.gov. The orbital wavefunctions of the target were obtained from atomic structure calculations using SUPERSTRUCTURE (Eissner et al. 1974), as in the earlier works of Bautista and Pradhan (1997) and Nahar et al. (1998). The present wavefunction expansion is a subset of the previous 31 term expansion, but includes singlet target terms which were not considered earlier. This enables the computation of the large number of doublet (N+1)-electron symmetries.

In addition to the target expansion in Table 1, it is necessary to ensure an accurate representation of short range correlation effects that are vital for the bound (N+1)-electron states. The second term of the wavefunction expansion, Eq.(1), consists of 61 such 'bound channel' correlation functions of Fe IV, as described by Bautista and Pradhan (1997).

The R-matrix calculations span several stages of computations (Berrington et al. 1987, 1995). The computations were extensive in terms of CPU memory size and time requirement. Hence, they were carried out one or a few symmetries of $SL\pi$ at a time. The one-electron orbital functions optimised through a configuration-interaction type calculation for Fe V using SUPERSTRUCTURE are the input to STG1 of the R-matrix codes to compute Slater integrals. Radial integrals for the partial wave expansion in Eq. 1 are specified for orbitals $0 \leq \ell \leq 9$ as a basis of NRANG2 = 12 'continuum' functions — sufficient for bound electrons with $n < 11$ at a radius of the R-matrix box. Computations are carried out for bound symmetries with $0 \leq L \leq 9$ (both parities) and spin multiplicities, $2S+1 = 2, 4, 6$.

The bound energy eigenvalues are obtained from computer program known STGB. All bound states of the Hamiltonian matrix are scanned for up to $n = 11$ and $l = 9$ with a fine effective quantum number mesh of $\Delta\nu = 0.001$. However, due

to the computational complexities stemming from the numerous bound states obtained, the code was unable to generate the standard R-matrix energy files. A complementary computer code, PLSRAD, was written to sort out the energies directly from the eigenvalues of the Hamiltonian matrix.

Furthermore, identification of the large number of LS bound states was a formidable task for this ion. With the exception of equivalent-electron states, the energies can be identified through quantum defect analysis of Rydberg series:

$$E = E_t - \frac{z^2}{\nu^2}, \quad \nu = n - \mu_{l\pm 1/2}(t) \quad (16)$$

with series limits E_t at the 16 Fe v ‘target’ states, and the weighted percentage contributions of interacting closed channels. An LS term is designated according to a possible combination of the configuration of the core and the outer electron, and assigned an appropriate value of ν for the outer electron quantum numbers n, l associated with the maximum channel percentage contribution. A Rydberg series is identified by sorting out the series of ν for the terms of same configuration but with increasing n of the valence electron. Code ELEVID (e.g. Nahar 1995) was used for this analysis. Uncertainties in assignments of proper configurations are introduced for cases with small differences in ν for various terms of the same symmetry, as the quantum defects for these states are almost the same. There is also some uncertainty in the identification of the adjacent higher or lower terms for such cases. The code STGBB is used to compute radiative data for the bound-bound transitions; the code exploits methods developed by Seaton (1986) to evaluate the outer region ($> R_A$) contributions to the radiative transition matrix elements.

The f-, S-, and A-values for the fine structure transitions are obtained through algebraic transformations of the LS multiplet line strengths using spectroscopically observed energies. For the cases where the observed energies are not found for all the fine structure components of an LS term, the spitting of LS f-values is carried out as described above using the code JYTOLS (Nahar 1995).

Using the new bound-bound radiative dataset, but retaining the earlier OP bound-free data (the contribution of bound-free is much smaller), we also calculate the monochromatic and Rosseland mean and Planck mean opacities as described by Seaton *et al.* (1994). Results are compared with those computed using the OP data calculated by Sawey and Berrington (1992).

4. Results and Discussion

In the subsections below we describe the results for bound state energies of LS terms and fine structure levels levels of Fe IV, radiative transitions for all dipole transitions, and the effect of the new data on opacities. We note that this entails the calculation and identification of the largest number of LS bound terms (1,798) computed for any ion under the Opacity or the Iron Projects thus far.

4.1. Energy Levels

A total of 1,798 bound states of Fe IV of sextet, quartet and doublet symmetries were obtained with $n \leq 11$, $l \leq 9$, $0 \leq L \leq 9$ (both even and odd parities), and spin multiplicities $2S+1 = 2, 4, 6$. All energy levels have been identified and a correspondence is established with the observed energies where available.

In Table 2, the present bound state energies are compared with the measured values (available from the NIST database), the OP calculations by Sawey and Berrington (1992), and by Bautista and Pradhan (1997). The low-lying energy levels in the present dataset are in very good agreement, within 5%, with the measured values. However, the higher ones show larger differences, the highest one being about 18% for the state $3d^4(^1D1)4p(^2D^o)$. A reason for such larger uncertainty for higher states, especially the odd parity states, could be the improper balance for inclusion of configurations which can improve the lower states considerably, but introduce some ‘spectral repulsion’ for the higher levels of the ion. Nonetheless, present energies on average are an improvement over the ones by Sawey and Berrington (1992) except for the high lying odd parity states where their energies show better agreement with experiment by a few percent. Present energies are also slightly better than those by Bautista and Pradhan (1997). The improvement in computed energies for the low-lying states is an important fact in the calculation of opacities, as discussed later.

The 104 observed LS term energies are obtained from statistical averaging of the 269 measured fine structure energies (NIST). In contrast, the present 1,798 LS states correspond to 5,331 fine structure levels; with the exception of the observed energy levels, the fine structure transition energies are approximated as explained in the Theory section.

The complete table of 1,798 bound state energies of Fe IV is available electronically. The energies of 104 observed states have been replaced by measured values. The bound states have been assigned with a prefixed index letter for convenience, as shown in sample Table 3 (the letters may not necessarily match with those of the NIST table). Following NIST, it is a standard practice to designate the lowest 52 bound states with prefixes in alphabetical order. For even symmetries, an ascending order of alphabet is used, first in the lower case and then in the upper case. The opposite order is chosen for the odd parity states, that is, descending order of alphabet, first in the lower case and then the upper case. States lying above these 52 states do not have any such notation; they are recognized by the energies only.

4.2. LS multiplets

The LS coupling oscillator strengths are used in various astrophysical applications, such as calculating plasma opacities. The LS multiplet data with 1,798 bound states of Fe IV comprises 138,121 dipole allowed oscillator strengths f and coefficients of spontaneous decay, the A – values. The full data table is available electronically. Since the line strength S does not depend on the transition energy, the present S values are used to recompute the f and A -values (Eq. 5) using observed energies whenever available to improve accuracy.

Table 2. Comparison of the present energies E_c of Fe IV with the observed binding energies E_o , and computed energies, E_{BP} (Bautista and Pradhan 1997) and E_{SB} (Sawey and Berrington 1992). An * next to a state indicates an incomplete multiplet with unobserved fine structure components.

<i>Config</i>	<i>S Lπ</i>	E_o (Ry)	E_c (Ry)	E_{BP} (Ry)	E_{SB} (Ry)
3d5	⁶ S	4.020000	3.978	3.984	3.96069
3d5	⁴ G	3.725831	3.658	3.665	3.64011
3d5	⁴ P	3.698270	3.652	3.660	3.58549
3d5	⁴ D	3.665794	3.607	3.613	3.56341
3d5	² I	3.590930	3.502		3.50204
3d5	³ ² D	3.566686	3.509		3.43429
3d5	² ² F	3.548645	3.494		3.42470
3d5	⁴ F	3.539584	3.478	3.485	3.42632
3d5	² H	3.507616	3.434		3.39455
3d5	² ² G	3.495274	3.421		3.38135
3d5	¹ ² F	3.462191	3.392		3.33754
3d5	² S	3.412002	3.348		3.27170
3d5	² ² D	3.344582	3.282		3.19871
3d5	¹ ² G	3.264596	3.186		3.12545
3d5	² P	3.107633	3.034		2.94101
3d5	¹ ² D	3.033568	2.965		2.85699
3d4(5D)4s	⁶ D	2.849206	2.872	2.941	2.77361
3d4(5D)4s	⁴ D	2.758817	2.778	2.795	2.68538
3d4(3P2)4s	⁴ P	2.607793	2.619	2.648	2.53137
3d4(3H)4s	⁴ H	2.612277	2.588	2.618	2.53129
3d4(3F2)4s	⁴ F	2.597263	2.592	2.613	2.51606
3d4(3G)4s	⁴ G	2.569849	2.549	2.580	2.48781
3d4(3P2)4s	² P	2.552383	2.561		2.47727
3d4(3H)4s	² H	2.556817	2.530		
3d4(3F2)4s	² F	2.543005	2.536		
3d4(3G)4s	² G	2.514627	2.492		2.43349
3d4(3D)4s	⁴ D	2.510927	2.494		2.42780
3d4(1G2)4s	² G	2.491359	2.476		2.40883
3d4(1I)4s	² I	2.484110	2.438		
3d4(1S2)4s	² S	2.464203	2.468		
3d4(3D)4s	² D	2.458112	2.440		2.37606
3d4(1D2)4s	² D	2.406789	2.388		2.32291
3d4(1F)4s	² F	2.350906	2.311		
3d4(5D)4p	⁶ F ^o	2.295457	2.219		2.25564
3d4(5D)4p	⁶ P ^o	2.287886	2.213		2.24990
3d4(3P1)4s	⁴ P	2.284212	2.246		2.19440
3d4(3F1)4s	⁴ F	2.285076	2.239		2.19842
3d4(5D)4p	⁴ P ^o	2.265726	2.183		2.22408
3d4(5D)4p	⁶ D ^o	2.258256	2.174		
3d4(3P1)4s	² P	2.232085	2.194		2.14451
3d4(3F1)4s	² F	2.232375	2.186		2.14682
3d4(5D)4p	⁴ F ^o	2.228839	2.142		2.18918
3d4(1G1)4s	² G	2.186593	2.125		2.09903
3d4(5D)4p	⁴ D ^o	2.176046	2.080		2.13458
3d4(3H)4p	⁴ H ^o	2.081972	1.977		2.03477
3d4(3P2)4p	⁴ D ^o	2.066669	1.945		2.02440
3d4(3F2)4p	⁴ G ^o	2.056962	1.940		2.00840
3d4(3H)4p	⁴ I ^o	2.050188	1.943		2.00550
3d4(3P2)4p	⁴ P ^o	2.039526	1.911		1.99660
3d4(3H)4p	² G ^o	2.049047	1.937		
3d4(3F2)4p	⁴ F ^o *	2.030249	1.907		1.98676
3d4(3H)4p	⁴ G ^o	2.031609	1.915		1.98801
3d4(3F2)4p	² D ^o *	2.025500	1.899		
3d4(3P2)4p	² P ^o	2.015051	1.889		
3d4(3H)4p	² I ^o	2.018812	1.905		
3d4(3F2)4p	⁴ D ^o	2.017690	1.887		1.96967

Table 3. Sample of energies in the complete table of energies.

<i>Config</i>	<i>S Lπ</i>	$E(Ry)$
3d5	a6S	-4.02000
3d4(5D)4d	b6S	-1.39828
3d4(5D)5d	c6S	-0.81636
3d4(5D)6d	d6S	-0.54091
3d4(5D)7d	e6S	-0.38568
3d4(5D)8d	f6S	-0.28891
3d4(5D)9d	g6S	-0.22449
3d4(5D)10d	h6S	-0.17945
3d4(5D)11d	i6S	-0.14672
3d4(5D)4d	a6P	-1.55722
3d4(5D)5d	b6P	-0.87599
3d4(5D)6d	c6P	-0.57173
3d4(5D)7d	d6P	-0.40367
3d4(5D)8d	e6P	-0.30030
3d4(5D)9d	f6P	-0.23216
3d4(5D)10d	g6P	-0.18486
3d4(5D)11d	h6P	-0.15068
...	...	...

Table 4 presents a sample of the complete electronic output file, with the following explanation. The first line provides the ion information, [Z=26, number of core electrons = 22]. Note that the number of core electrons is one less than that in the final ion. This (somewhat confusing) notation is standard in the R-matrix calculations and refers to the ‘target ion’ with which another electron forms a bound state. The first two columns, [SiLiPi(i) -> SflfPf(f)], refers to the symmetries of the initial and final bound states; the next two columns to the initial and final energies, E_i and E_f , in Rydbergs, followed by f , gf , A -values. The total number of transitions is (138121) is also listed at the end of the headings. The A -values are in atomic units; they should be divided by 2.419×10^{-17} to yield $A(sec^{-1})$. Designation for a given state in the first two columns consists of integers representing the spin multiplicity ‘S’, which is in fact (2S+1); the total angular momentum L is 0 for S, 1 for P, etc.; the parity $P_i = 0$ for an even state and = 1 for an odd state; and the fourth integer is the index of the ascending energy order computed by STGB for the given symmetry.

Present results for the Fe IV oscillator strengths are compared with earlier works in Table 5. Bautista and Pradhan (1997) considered transitions among sextets and quartets only and hence have relatively smaller set of data compared to the OP data. Under the OP, Sawey and Berrington (1992) also produced an extensive set of data, a total of 133,650 transitions among 1,721 bound states for all three symmetries, sextets, quartets, and doublets. Hence, OP data contains about 4,500 transitions less than the present set. Fawcett (1989) employed the semi-empirical Cowan’s Hartree-Fock relativistic code with optimized Slater parameters to calculate the transition probabilities of Fe IV. We note that the NIST database of evaluated compilation gives radiative decay rates only for a small number of forbidden transitions, not allowed ones. Table 5 compares present results with these earlier works for some transitions. Agreement among all the calculations is good for sextet and quartet transitions, within 10%. However, for the doublet tran-

sitions the present oscillator strengths and those of Sawey and Berrington (1992) show similar values, while Fawcett's values are somewhat smaller.

4.3. Fine structure transitions

Fine structure f - and A -values are presented for 712,120 dipole allowed transitions in Fe IV. As in the case of LS multiplets, observed energies of fine structure transitions are employed whenever available for improved accuracy. As only a limited number of observed levels are available, a smaller set of 2,915 fine structure transitions corresponding to 620 LS multiplets has observed energies. Rest of the transitions, 709,205 in total, employ calculated energies; the transition energies are obtained as explained in the Theory section. The entire file containing all fine structure transitions is available electronically.

A sample set of f -, S -, and A -values for the dipole allowed fine structure transitions in Fe IV is presented in Table 6. The first line of each subset corresponds to the LS transition followed by the fine structure components. The letter prefix designation of the transitional states in the table correspond to their energy positions, as explained above (see Table 3). The energies of the initial and final fine structure levels are given in unit of cm^{-1} , while the transitional energy differences are given in Rydbergs. The A -values are in s^{-1} . An asterisk (*) below an LS state indicates an incomplete set of observed energy levels, and an asterisk for the transitional energy indicates that one or both the levels are missing from the observed energy set.

Based on the accuracy of the calculated energies and comparison with other works, LS multiplets should be of better accuracy than those in TOPbase and by Bautista and Pradhan (1997). The overall accuracy for most of the present f -, S -, and A -values should be within 20% for strong transitions, with the possible exception of weak transitions. Because of low charge of the ion, the relativistic effects are expected to be small.

4.4. Opacities

We calculate sample radiative opacities of Fe IV for three reasons. First, as a completeness and consistency check on the huge amount of data computed, second, to gauge the potential impact on iron opacities, and third, spectral models for determination of iron abundances in astrophysical sources.

Fig. 1 shows the monochromatic opacities in most of the wavelength range where Fe IV is a contributor in an astrophysical plasma in local-thermodynamic-equilibrium (LTE). The present results are compared with those using the bound-bound OP data by Sawey and Berrington (1992). In order to compare only the main source of opacity (lines), we have used the same bound-free (photoionization) data in both calculations; the bound-free opacity is less than 10% of the total. We choose the temperature $\log(T) = 4.5$ and the electron density $N_e = 17.0$. The Mihalas-Hummer-Dappen equation-of-state (MHD-EOS; Mihalas *et al.* 1998) used in the OP work gives relatively large value of the ionization fraction $\text{Fe IV}/\text{Fe} = 0.57$; Fe IV dominates the iron opacity at this temperature and density. Fig. 1 shows immediately the reason that Fe IV is the dominant ion in

the UV, between 500-3000 Å. It is noteworthy that the opacity of Fe IV varies over nearly 10 orders of magnitude, implying that it remains a large contributor over a very wide range of radiation.

The two datasets displayed in Fig. 1 differ considerably. The main features throughout the opacity spectrum appear at different positions. For example, there is relative displacement of nearly 200 Å in the position of the trough which is at ~ 1500 Å in the present results, but at ~ 1700 Å using the OP data. Even more striking are the differences in the present and the OP Rosseland mean opacities, 3.3 and $4.87 \text{ cm}^2/\text{g}$ respectively. This nearly 50% difference is similar to the one we found earlier in the Fe II opacities (Nahar and Pradhan 1994) using new IP data as opposed to the OP data by Sawey and Berrington (1997), which has been shown to be inaccurate for low ionization states of iron Fe I- IV (Nahar and Pradhan 1994, Bautista and Pradhan 1997, Cowley and Bautista 2003). We note that the calculation of latest stellar opacities under the OP has been carried out using the OP data as well as data from other sources (M.J. Seaton, private communication).

Although more extensive investigations are needed to ascertain the full range of differences at all temperatures and densities, we analyze the results somewhat further. The improvement in computed energies in the present calculations for the low-lying states is an important fact in the calculation of opacities, since they account for most of the level population; higher levels are (a) not populated significantly, and (b) smeared out or dissolved due to Stark and electron impact broadening as incorporated in MHD-EOS. An examination of the all level populations of Fe IV at $T = 3.16 \times 10^5 \text{ K}$ and $N_e = 10^{17} \text{ cm}^{-3}$, obtained from the MHD-EOS shows that there is nearly a sharp drop in population of levels as one approaches the highest of the 16 ground configuration $3d^5$ levels. For example, the population drops by about 2 orders of magnitude from the ground state $3d^5(^6S)$ at $\sim -4 \text{ Ryd}$, to the first of the excited configuration state $3d^44s(^6D)$ at $\sim -2.8 \text{ Ryd}$. Even higher levels do not contribute to the opacity in any significant way. The decrease is related to the negative exponential functional form in the modified Boltzmann-Saha equations in MHD-EOS.

5. Conclusions

Results are presented from a large and extensive calculation for Fe IV. All 1,798 states were identified in spectroscopic notation through quantum defect and channel contribution analysis. However, level identifications for highly mixed excited states should be treated with caution since often no definitive assignment is possible, and in any case not needed for practical applications. Present energies agree very well with the observed energies for all low lying terms. However, the uncertainty increases for excited states with higher energies, especially the odd parity states. A small subset of the total number of corresponding 138,121 transitions among the LS terms, and 712,120 fine structure transitions, was reprocessed wherever observed energies are available. While the transition probabilities for the strong transitions should not exceed about 20%, the uncertainty in the weak fine structure components could be

much higher, especially if relativistic effects are important and deviations from LS coupling are significant.

The new Fe IV results should be particularly useful for the analysis and application to ultraviolet to optical spectra from astrophysical and laboratory sources. All data files are electronically available from the CDS.

Acknowledgements. This work was partially supported by NASA (SNN) and U.S. National Science Foundation (AKP). The computational work was largely carried out on the Cray SV1 at the Ohio Supercomputer Center in Columbus, Ohio.

References

- Allen, C.W., 1976, *Astrophysical Quantities* (3rd Edition, Athlone Press, London)
- Bautista, M.A. & Pradhan, A.K., 1997, *Astron. Astrophys. Suppl.* 126, 365
- Berrington, K.A., Burke, P.G., Butler, K., Seaton, M.J., Storey, P.J., Taylor, K.T., Yu, Y., 1987, *J. Phys. B* 20, 6379
- Berrington, K.A., Eissner, W.B. & Norrington, P.H. 1995, *Comput. Phys. Commun* 92, 290
- Cowley, C.R. and Bautista, M.A., 2003, *Mon. Not. R. Astro. Soc.* 341, 1226
- Cunto, W., Mendoza, C., Ochsenbein, F. & Zeippen, C.J., 1993, *A&A* 275, L5 (TOPbase). The website addresses are <http://vizier.u-strasbg.fr/OP.html> and <http://heasarc.gsfc.nasa.gov>
- Dopita, M.A. and Sutherland, R.S., 2003 *Astrophysics of the Diffuse Universe*, Springer-Verlag.
- Eissner, W, Jones, M. & Nussbaumer, H, 1974, *Comput. Phys. Commun* 8, 270
- Eissner, W. & Zeippen, C.J., 1981 *J. Phys. B* 14, 2125
- Fawcett, B.C., 1989, *At. Data Nucl. Data Tables* 41, 181
- Garstang, R.H., 1958, *Mon. Not. R. Astro. Soc.* 118, 572
- Hubeny, I., Mihalas, D. & Werner, K., 2003, *Stellar Atmospheres Modeling*, ASP Conference Proceedings, Vol. 288 (ISBN:1-58381-131-1)
- Hyung, S. and Aller, L.H. 1998, *PASP* 110, 466
- Hummer, D.G, Berrington, K.A., Eissner, W, Pradhan, A.K, Saraph, H.E., & Tully, J.A., 1993, *A&A* 279, 298 **IP**
- Mihalas, D., Dappen, W. and Hummer, D.G., 1988, *ApJ* 331, 815
- McKenna, F.C., Keenan, F.P., Hambly, N.C., Allende, P.C., Rolleston, W.R., J., Aller, L.H., Feibelman, W.A., 1997, *ApJS* 109, 225
- Nahar, S.N., 1993, *Physica Scr.* 48, 297
- Nahar, S.N., 1995, *A&A* 293, 967
- Nahar, S.N., 1997, *Physica Scr.* 55, 200
- Nahar, S.N., Bautista, M.A., & Pradhan A.K., 1998, *Phys. Rev. A* 58, 4593
- Nahar, S.N., Delahaye, F., Pradhan, A.K., & Zeippen, C.J., 2000, *Astron. Astrophys. Suppl. Ser.* 144, 141
- Nahar, S.N. & Pradhan, A.K., 1996, *A&AS* 119, 509
- Nahar, S.N. & Pradhan, A.K., 2000, *Phys. Scr.* 61, 675
- Nahar, S.N., Werner, W., Chen, G.X., & Pradhan, A.K., 2003, *A&A* 408, 789
- NIST: www.nist.gov
- Rubin, R. *et al.* 1997, *ApJ* 474, L131
- Rubin, R. and Froese Fisher, C. 1998, *J. Phys. B* 31, 1657
- Sawey, P.M. & Berrington, K.A., 1992, *J. Phys. B* 25, 1451
- Seaton, M.J., 1985, *J. Phys. B* 18, 2111
- Seaton, M.J., 1986, *J. Phys. B* 19, 2601
- Seaton, M.J., 1987, *J. Phys. B* 20, 6363
- Seaton, M.J., Yan, Yu., Mihalas, D., & Pradhan, A.K., 1994, *Mon. Not. R. Astron. Soc.* 266, 805
- Sigut, T.A.A. & Pradhan, A.K. 2003, *ApJS* 100, 100
- Sigut, T.A.A., Pradhan, A.K., Nahar, S.N. 2004 (in press, *astro-ph/0401168*)
- The Opacity Project vols 1 & 2*, compiled by the Opacity Project Team: Institute of Physics, London UK 1995 and 1996 — **OP**.

Table 4. Oscillator strengths, f , gf -values and radiative decay rates, A for transitions in Fe IV [$Z=26$, number of core electrons=22]. The first transition, [6 0 0 1 6 1 1 1], means ${}^6S(1) \rightarrow {}^6P^o(1)$, where the number within parenthesis is the energy position of the state. The energies are Rydberg and the A -values are in atomic units, i.e. $A(\text{sec}^{-1}) = A(a.u.)/2.419 \times 10^{-17}$.

26	22									
SiLiPi(i) ->	SfLfPf(f)	Ei	Ef	fif	gfl	Afi				
138121										
6 0 0 1	6 1 1 1	-4.020000	-2.287886	4.2259E-01	2.536E+00	3.395E+09				
6 0 0 1	6 1 1 2	-4.020000	-1.163950	4.8711E-02	2.923E-01	1.064E+09				
6 0 0 1	6 1 1 3	-4.020000	-1.017630	4.3453E-01	2.607E+00	1.049E+10				
6 0 0 1	6 1 1 4	-4.020000	-0.717120	1.7706E-02	1.062E-01	5.172E+08				
6 0 0 1	6 1 1 5	-4.020000	-0.650580	2.3136E-01	1.388E+00	7.032E+09				
6 0 0 1	6 1 1 6	-4.020000	-0.487200	8.8916E-03	5.335E-02	2.971E+08				
6 0 0 1	6 1 1 7	-4.020000	-0.451150	1.2883E-01	7.730E-01	4.393E+09				
6 0 0 1	6 1 1 8	-4.020000	-0.352670	5.1805E-03	3.108E-02	1.865E+08				
6 0 0 1	6 1 1 9	-4.020000	-0.331080	7.7609E-02	4.657E-01	2.828E+09				
6 0 0 1	6 1 1 10	-4.020000	-0.267150	3.3135E-03	1.988E-02	1.249E+08				
6 0 0 1	6 1 1 11	-4.020000	-0.253220	4.9970E-02	2.998E-01	1.898E+09				
6 0 0 1	6 1 1 12	-4.020000	-0.209400	2.2580E-03	1.355E-02	8.778E+07				
6 0 0 1	6 1 1 13	-4.020000	-0.199880	3.3953E-02	2.037E-01	1.327E+09				
6 0 0 1	6 1 1 14	-4.020000	-0.168560	1.6124E-03	9.675E-03	6.404E+07				
6 0 0 1	6 1 1 15	-4.020000	-0.161760	2.4110E-02	1.447E-01	9.609E+08				
6 0 0 1	6 1 1 16	-4.020000	-0.138610	1.1931E-03	7.159E-03	4.812E+07				
6 0 0 1	6 1 1 17	-4.020000	-0.133580	1.7723E-02	1.063E-01	7.167E+08				
6 1 1 1	6 0 0 2	-2.287886	-1.398280	2.1035E-01	3.786E+00	4.011E+09				
6 0 0 2	6 1 1 2	-1.398280	-1.163950	3.2583E-01	1.955E+00	4.792E+07				
6 0 0 2	6 1 1 3	-1.398280	-1.017630	1.0472E+00	6.283E+00	4.062E+08				
6 0 0 2	6 1 1 4	-1.398280	-0.717120	5.6950E-03	3.417E-02	7.075E+06				
6 0 0 2	6 1 1 5	-1.398280	-0.650580	3.3867E-03	2.032E-02	5.069E+06				
6 0 0 2	6 1 1 6	-1.398280	-0.487200	1.4603E-03	8.762E-03	3.246E+06				
6 0 0 2	6 1 1 7	-1.398280	-0.451150	1.1753E-02	7.052E-02	2.823E+07				
6 0 0 2	6 1 1 8	-1.398280	-0.352670	6.6550E-04	3.993E-03	1.948E+06				
6 0 0 2	6 1 1 9	-1.398280	-0.331080	9.9583E-03	5.975E-02	3.037E+07				
6 0 0 2	6 1 1 10	-1.398280	-0.267150	3.7717E-04	2.263E-03	1.292E+06				
6 0 0 2	6 1 1 11	-1.398280	-0.253220	7.4100E-03	4.446E-02	2.601E+07				
6 0 0 2	6 1 1 12	-1.398280	-0.209400	2.3900E-04	1.434E-03	9.045E+05				
6 0 0 2	6 1 1 13	-1.398280	-0.199880	5.4667E-03	3.280E-02	2.102E+07				
6 0 0 2	6 1 1 14	-1.398280	-0.168560	1.6250E-04	9.750E-04	6.579E+05				
6 0 0 2	6 1 1 15	-1.398280	-0.161760	4.0933E-03	2.456E-02	1.676E+07				
6 0 0 2	6 1 1 16	-1.398280	-0.138610	1.1605E-04	6.963E-04	4.930E+05				
6 0 0 2	6 1 1 17	-1.398280	-0.133580	3.1250E-03	1.875E-02	1.339E+07				
.....										

Table 5. Comparison of present f -values for Fe IV with previous works: a - Bautista and Pradhan (1997), b - Sawey and Berrington (1992), c - Fawcett (1989).

Present	Others	$C_i - C_k$	$SL_i - SL_k$	$J_i - J_k$
0.1817	0.2037 ^a , 0.1973 ^b , 0.2037 ^c	$3d^4({}^5D)4s - 3d^4({}^5D)4p$	${}^6D - {}^6P^o$	30 - 18
0.3085	0.339 ^a , 0.3277 ^b , 0.2830 ^c	$3d^4({}^5D)4s - 3d^4({}^5D)4p$	${}^6D - {}^6D^o$	30 - 30
0.4142	0.4603 ^a , 0.4433 ^b , 0.4633 ^c	$3d^4({}^5D)4s - 3d^4({}^5D)4p$	${}^6D - {}^6F^o$	30 - 42
0.1710	0.1883 ^a , 0.1820 ^b , 0.1100 ^c	$3d^4({}^5D)4s - 3d^4({}^5D)4p$	${}^4D - {}^4P^o$	20 - 12
0.3165	0.351 ^a , 0.3335 ^b , 0.3285 ^c	$3d^4({}^5D)4s - 3d^4({}^5D)4p$	${}^4D - {}^4D^o$	20 - 20
0.4164	0.4583 ^a , 0.4395 ^b , 0.4375 ^c	$3d^4({}^5D)4s - 3d^4({}^5D)4p$	${}^4D - {}^4F^o$	30 - 28
0.1599	0.1491 ^b , 0.1073 ^c	$3d^4({}^5H)4s - 3d^4({}^5H)4p$	${}^2H - {}^2G^o$	22 - 18
0.2679	0.2559 ^b , 0.2018 ^c	$3d^4({}^5H)4s - 3d^4({}^5H)4p$	${}^2H - {}^2H^o$	22 - 22
0.3472	0.3645 ^b , 0.3136 ^c	$3d^4({}^5H)4s - 3d^4({}^5H)4p$	${}^2H - {}^2I^o$	22 - 26




Fig. 1. Monochromatic opacities of Fe IV using the present data (top panel) and with the Opacity Project data (lower panel) from Sawey and Berrington (1992) at $\text{Log } T \text{ (K)} = 4.5$ and $\text{Log } N_e \text{ (cm}^{-3}\text{)} = 17.0$. The major features show a shift in wavelength or differences in intensities.

Table 6. Sample data for allowed fine structure transitions of Fe IV.

$SL_i - SL_k$	E_i Ry/cm^{-1}	E_k Ry/cm^{-1}	E_{ik} $\text{Ry}/\text{\AA}$	g_i	g_k	f_{ik}	S	A s^{-1}
$a\ ^6S^e \rightarrow z\ ^6P^o$	4.0200	2.2879	1.732E+00	6	18	4.226E-01	4.392E+00	3.395E+09
	0.000	190226.00	5.257E+01	6	8	1.880E-01	1.952E+00	3.403E+09
	0.000	190008.00	5.263E+01	6	6	1.408E-01	1.464E+00	3.391E+09
	0.000	189885.00	5.266E+01	6	4	9.381E-02	9.759E-01	3.384E+09
$a\ ^6D^e \rightarrow z\ ^6P^o$	2.8492	2.2879	5.613E-01	30	18	1.827E-01	2.929E+01	7.704E+08
	128967.000	190226.00	1.632E+02	10	8	1.817E-01	9.762E+00	5.684E+08
	128541.000	190226.00	1.621E+02	8	8	6.554E-02	2.798E+00	1.664E+08
	128191.000	190226.00	1.612E+02	6	8	1.461E-02	4.653E-01	2.813E+07
	128541.000	190008.00	1.627E+02	8	6	1.170E-01	5.011E+00	3.930E+08
	128191.000	190008.00	1.618E+02	6	6	1.120E-01	3.579E+00	2.855E+08
	127929.000	190008.00	1.611E+02	4	6	5.523E-02	1.171E+00	9.464E+07
	128191.000	189885.00	1.621E+02	6	4	5.692E-02	1.822E+00	2.167E+08
	127929.000	189885.00	1.614E+02	4	4	1.286E-01	2.733E+00	3.293E+08
	127766.000	189885.00	1.610E+02	2	4	1.842E-01	1.952E+00	2.371E+08
	2.8492	2.2583	5.910E-01	30	30	3.085E-01	4.698E+01	8.654E+08
	128967.000	193789.00	1.543E+02	10	10	2.510E-01	1.275E+01	7.035E+08
$a\ ^6D^e \rightarrow z\ ^6D^o$	128967.000	193386.00	1.552E+02	10	8	5.669E-02	2.897E+00	1.962E+08
	128541.000	193789.00	1.533E+02	8	10	7.178E-02	2.897E+00	1.631E+08
	128541.000	193386.00	1.542E+02	8	8	1.380E-01	5.607E+00	3.872E+08
	128541.000	192595.00	1.561E+02	8	6	9.789E-02	4.025E+00	3.572E+08
	128541.000	192595.00	1.561E+02	8	6	9.789E-02	4.025E+00	3.572E+08
	128191.000	193386.00	1.534E+02	6	8	1.328E-01	4.025E+00	2.825E+08
	128191.000	192595.00	1.553E+02	6	6	5.249E-02	1.610E+00	1.452E+08
	128191.000	193271.00	1.537E+02	6	4	1.238E-01	3.759E+00	5.248E+08
	127929.000	192595.00	1.546E+02	4	6	1.846E-01	3.759E+00	3.432E+08
	127929.000	193271.00	1.530E+02	4	4	3.419E-03	6.891E-02	9.738E+06
	127929.000	193120.00	1.534E+02	4	2	1.210E-01	2.443E+00	6.857E+08
	127766.000	193271.00	1.527E+02	2	4	2.431E-01	2.443E+00	3.478E+08
	127766.000	193120.00	1.530E+02	2	2	6.933E-02	6.985E-01	1.975E+08
	2.8492	2.2955	5.537E-01	30	42	4.142E-01	6.731E+01	7.286E+08
	128967.000	190276.00	1.631E+02	10	12	3.582E-01	1.923E+01	7.483E+08
	128967.000	189515.00	1.652E+02	10	10	5.424E-02	2.949E+00	1.326E+08
	128967.000	188904.00	1.668E+02	10	8	4.319E-03	2.372E-01	1.293E+07
	128541.000	189515.00	1.640E+02	8	10	3.020E-01	1.305E+01	5.992E+08
$a\ ^6D^e \rightarrow z\ ^6F^o$	128541.000	188904.00	1.657E+02	8	8	9.991E-02	4.359E+00	2.428E+08
	128541.000	188428.00	1.670E+02	8	6	1.246E-02	5.481E-01	3.975E+07
	128191.000	188904.00	1.647E+02	6	8	2.542E-01	8.270E+00	4.687E+08
	128191.000	188428.00	1.660E+02	6	6	1.359E-01	4.456E+00	3.288E+08
	128191.000	188086.00	1.670E+02	6	4	2.333E-02	7.693E-01	8.373E+07
	127929.000	188428.00	1.653E+02	4	6	2.121E-01	4.616E+00	3.451E+08
	127929.000	188086.00	1.662E+02	4	4	1.669E-01	3.654E+00	4.029E+08
	127929.000	187878.00	1.668E+02	4	2	3.240E-02	7.116E-01	1.553E+08
	127766.000	188086.00	1.658E+02	2	4	1.821E-01	1.987E+00	2.209E+08
	127766.000	187878.00	1.664E+02	2	2	2.283E-01	2.500E+00	5.502E+08
	3.6983	1.6735	2.025E+00	12	4	6.310E-02	1.122E+00	6.233E+09
	35253.800	257503.00	4.499E+01	6	4	6.311E-02	5.609E-01	3.119E+09
$a\ ^4P^e \rightarrow z\ ^4S^o$	35333.300	257503.00	4.501E+01	4	4	6.309E-02	3.739E-01	2.077E+09
	35406.600	257503.00	4.503E+01	2	4	6.307E-02	1.870E-01	1.038E+09
	3.6983	2.2657	1.433E+00	12	12	5.704E-02	1.433E+00	9.402E+08
	35253.800	193549.00	6.317E+01	6	6	4.020E-02	5.017E-01	6.720E+08
$a\ ^4P^e \rightarrow z\ ^4P^o$	35253.800	191694.00	6.392E+01	6	4	1.703E-02	2.150E-01	4.170E+08
	35333.300	193549.00	6.320E+01	4	6	2.583E-02	2.150E-01	2.875E+08
	35333.300	191694.00	6.395E+01	4	4	7.564E-03	6.371E-02	1.234E+08
	35333.300	191021.00	6.423E+01	4	2	2.354E-02	1.991E-01	7.611E+08
	35406.600	191694.00	6.398E+01	2	4	4.726E-02	1.991E-01	3.849E+08
	35406.600	191021.00	6.426E+01	2	2	9.410E-03	3.982E-02	1.520E+08